arXiv:math/0505133v1 [math.NT] 8 May 2005

MULTIVARIATE p-ADIC L-FUNCTION

Taekyun Kim

Institute of Science Education,
Kongju National University, Kongju 314-701, S. Korea
e-mail: tkim64@hanmail.net (or tkim@kongju.ac.kr)

Abstract. In the recent, many mathematicians studied the multiple zeta function in the complex number field. In this paper we construct the p-adic analogue of multiple zeta function which interpolates the generalized multiple Bernoulli numbers attached to χ at negative integers.

§1. Introduction

Let p be a fixed prime. Throughout this paper $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$ will, respectively, denote the ring of p-adic rational integers, the field of p-adic rational numbers, the complex number field and the completion of algebraic closure of $\mathbb{Q}_p$, cf.[3, 10, 17]. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-v_p(p)} = p^{-1}$. The Bernoulli numbers in $\mathbb{C}$ are defined by

$$F(t) = \frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!}, \text{ for } |t| < 2\pi. \tag{1}$$

From the definition, one has $B_0 = 1, B_1 = -\frac{1}{2}, \cdots$. Also $B_{2k+1} = 0$ for $k \geq 1$. Bernoulli numbers are used to express the special values of Riemann zeta function $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$, for $s \in \mathbb{C}$, namely $\zeta(2m) = \frac{(2\pi)^{2m}(-1)^{m-1}B_{2m}}{2(2m)!}$, $m \geq 1$. For the

---

Key words and phrases. p-adic q-integrals, multiple Barnes' Bernoulli numbers.
2000 Mathematics Subject Classification: 11S80, 11B68, 11M99 .

Typeset by $\mathcal{AMS}$-TEX

negative integers, we note that $\zeta(1-2m) = -\frac{B_{2m}}{2m}$. The multiple Bernoulli numbers of order r are defined as

$$F^r(t) = \left(\frac{t}{e^t-1}\right)^r = \sum_{n=0}^{\infty} B_n^{(r)} \frac{t^n}{n!}, \text{ for } |t| < 2\pi,\ r \in \mathbb{N}. \tag{2}$$

Let x be an indeterminate. Then the multiple Bernoulli polynomials are also defined by

$$F^r(t,x) = \left(\frac{t}{e^t-1}\right)^r e^{xt} = \sum_{n=0}^{\infty} B_n^{(r)}(x) \frac{t^n}{n!}, \text{ for } |t| < 2\pi,\ r \in \mathbb{N}. \tag{3}$$

Let χ be a primitive Dirichlet character with conductor $f \in \mathbb{N}$. The generalized Bernoulli numbers attached to χ, $B_{n,\chi}$, are defined by

$$F_\chi(t) = \sum_{a=1}^{f} \frac{\chi(a)te^{at}}{e^{ft}-1} = \sum_{n=0}^{\infty} B_{n,\chi} \frac{t^n}{n!}, \text{ for } |t| < \frac{2\pi}{f}\ . \tag{4}$$

In [8, 9], the multiple generalized Bernoulli numbers attached to χ, $B_{n,\chi}^{(r)}$, are defined by

$$F_\chi^r(t) = \sum_{a_1,\cdots,a_r=1}^{f} \frac{\chi(\sum_{i=1}^{r} a_i) t^r e^{t\sum_{i=1}^{r} a_i}}{(e^{ft}-1)^r} = \sum_{n=0}^{\infty} B_{n,\chi}^{(r)} \frac{t^n}{n!},\ |t| < \frac{2\pi}{f}\ . \tag{5}$$

Kubota and Leopoldt proved the existence of meromorphic functions, $L_p(s,\chi)$, defined over the p-adic number field, that serve as p-adic equivalents of the Dirichlet L-series. These p-adic L-functions interpolate the values

$$L_p(1-n,\chi) = -\frac{1}{n}(1-\chi_n(p)p^{n-1})B_{n,\chi_n}, \text{ for } n \in \mathbb{N} = \{1,2,\cdots,\}\ ,$$

where $B_{n,\chi}$ denote the nth generalized Bernoulli numbers associated with the primitive Dirichlet character χ, and $\chi_n = \chi w^{-n}$, with w the *Teichmüller* character, cf.[1-24]. In this paper we study analytic continued function which interpolates the multiple generalized Bernoulli numbers attached to χ at negative integers in complex plane. Finally we will construct multivariate p-adic L-function by using Washington's method.

2. MULTIVARIATE DIRICHLET'S L-FUNCTION ASSOCIATED WITH THE MULTIPLE GENERALIZED BERNOULLI NUMBERS ATTACHED TO χ AT NEGATIVE INTEGERS IN $\mathbb{C}$

In [8, 9], the multiple generalized numbers attached to χ, $B_{n,\chi}^{(r)}$, are defined by

$$F_{\chi}^{r}(t)=\sum_{a_1,\cdots,a_r=1}^{f}\frac{\chi(\sum_{i=1}^{r}a_i)t^re^{t\sum_{i=1}^{r}a_i}}{(e^{ft}-1)^r}=\sum_{n=0}^{\infty}B_{n,\chi}^{(r)}\frac{t^n}{n!},\ |t|<\frac{2\pi}{f}\ .$$

By (3) and (5), we easily see that

$$B_{n,\chi}^{(r)}=f^{n-r}\sum_{a_1,\cdots,a_r=0}^{f-1}B_n^{(r)}(\frac{a_1+\cdots+a_r}{f})\chi(a_1+\cdots+a_r),\ \text{cf. [8, 9]}\ . \tag{6}$$

For $s\in\mathbb{C}$, multiple Hurwitz's zeta function is defined by

$$\zeta_r(s,x)=\sum_{n_1,\cdots,n_r=0}\frac{1}{(x+n_1+\cdots+n_r)^s}=\frac{1}{\Gamma(s)}\int_0^{\infty}F_r(-t,x)t^{s-r-1}dt,\ \text{cf. [7, 8]}\ . \tag{7}$$

Note that

$$\zeta_r(-n,x)=(-1)^r\frac{n!}{(n+r)!}B_{n+r}^{(r)}(x),\ \text{for } n\in\mathbb{N},\ \text{cf. [16]}\ .$$

We also consider the below complex integral in $\mathbb{C}$:

$$\frac{1}{\Gamma(s)}\int_0^{\infty}F_{\chi}^{r}(-t)t^{s-r-1}dt=\sum_{\substack{n_1,\cdots,n_r=0\\ n_1+\cdots+n_r\neq0}}^{\infty}\frac{\chi(n_1+\cdots+n_r)}{(n_1+\cdots+n_r)^s},\ \text{for } s\in\mathbb{C},\ |t|<\frac{2\pi}{f}, \tag{8}$$

where χ is the primitive Dirichlet's character with conductor $f\in\mathbb{N}$.

From Eq.(8), we can derive the multivariate Dirichlet's L-function in complex plane as follows:

Definition 1. *For $s\in\mathbb{C}$, define*

$$L_r(s,\chi)=\sum_{\substack{n_1,\cdots,n_r=0\\ n_1+\cdots+n_r\neq0}}^{\infty}\frac{\chi(n_1+\cdots+n_r)}{(n_1+\cdots+n_r)^s}, \tag{9}$$

where χ is the Dirichlet's character with conductor $f\in\mathbb{N}$.

Note that $L_r(s,\chi)$ is meromorphic for $s\in\mathbb{C}$ with poles at $s=1,2,\cdots,r$. By (8) and (5), we obtain the following:

Theorem 2. *For $n \in \mathbb{N}$, we have*

$$L_r(-n,\chi) = (-1)^r \frac{n!}{(n+r)!} B^{(r)}_{n+r,\chi}. \tag{10}$$

Let s be a complex variable, a, F be integers with $0 < a < F$. Then we now consider the function $H_r(s; a_1, \cdots, a_r | F)$ as follows:

$$H_r(s; a_1, \cdots, a_r|F) = \sum_{\substack{m_1,\cdots,m_r>0 \\ m_i \equiv a_i (\mod F)}} \frac{1}{(m_1+\cdots+m_r)^s} = F^{-s}\zeta_r(s, \frac{a_1+\cdots+a_r}{F}). \tag{11}$$

The function $H_r(s; a_1, \cdots, a_r|F)$ is a meromorphic in whole complex plane with poles at $s = 1, 2, \cdots, r$.

Let $\chi(\neq 1)$ be the Dirichlet's character with conductor $F \in \mathbb{N}$. Then the multivariate Dirichlet's L-function can be expressed as the sum

$$L_r(s,\chi) = \sum_{a_1,\cdots,a_r=1}^{F} \chi(a_1+\cdots+a_r) H_r(s; a_1, \cdots, a_r|F) \text{ for } s \in \mathbb{C}\ . \tag{12}$$

By simple calculation, we easily see that

$$H_r(-n; a_1, \cdots, a_r|F) = F^n (-1)^r \frac{n!}{(n+r)!} B^{(r)}_{n+r}(\frac{a_1+\cdots+a_r}{F}) \text{ for } r, n \in \mathbb{N}. \tag{13}$$

Thus, we have

$$L_r(-n,\chi) = (-1)^r \frac{n!}{(n+r)!} B^{(r)}_{n+r,\chi} \text{ for } n \in \mathbb{N}\ . \tag{14}$$

By using Eq.(3), the Eq.(13) is modified by

$$H_r(s; a_1, \cdots, a_r|F) = \frac{1}{F^r} \frac{(a_1+\cdots+a_r)^{-s+r}}{\prod_{j=1}^r (s-j)} \sum_{k=0}^{\infty} \binom{-s+r}{k} \left(\frac{F}{a_1+\cdots+a_r}\right)^k B^{(r)}_k. \tag{15}$$

From (12),(13) and (15), we can derive the below:

$$\begin{aligned} L_r(s,\chi) = &\frac{1}{\prod_{j=1}^r (s-j)} \frac{1}{F^r} \sum_{a_1,\cdots,a_r=1}^{F} \chi(a_1+\cdots+a_r)(a_1+\cdots+a_r)^{r-s} \\ &\cdot \sum_{m=0}^{\infty} \binom{r-s}{m} \left(\frac{a_1+\cdots+a_r}{F}\right)^m B^{(r)}_m. \end{aligned} \tag{16}$$

Remark. The values of $L_r(s,\chi)$ at negative integers are algebraic, hence may be regarded as lying in an extension of $\mathbb{Q}_p$. We therefore look for a p-adic function which agrees with $L_r(s,\chi)$ at negative integers in the next section.

3. Multivariate p-adic L-function

In this section we shall consider the p-adic analogs of the multivariate L-function, $L_r(s,\chi)$, which were introduced in the previous section. Indeed this function is the p-adic interpolation function for the generalized multiple Bernoulli numbers attached to χ. Let w denote the *Teichmüller* character, having conductor $f_w = q$. For an arbitrary character χ, we define $\chi_n = \chi w^{-n}$, where $n \in \mathbb{Z}$, in the sense of the product of characters. Let $< a >= w^{-1}(a)a = \frac{a}{w(a)}$. Then, we note that $< a > \equiv 1 \pmod{q\mathbb{Z}_p}$. Let $A_j(x) = \sum_{n=0}^{\infty} a_{n,j}x^n$, $a_{n,j} \in \mathbb{C}_p$, $j = 0,1,2,\cdots$ be a sequence of power series, each of which converges in a fixed subset $D = \{s \in \mathbb{C}_p \,|\, |s|_p \le |p^*|^{-1} p^{-\frac{1}{p-1}}\}$ of $\mathbb{C}_p$ such that (1) $a_{n,j} \to a_{n,0}$ as $j \to \infty$ for $\forall n$; (2) for each $s \in D$ and $\epsilon > 0$, there exists $n_0 = n_0(s,\epsilon)$ such that $\left|\sum_{n\ge n_0} a_{n,j}s^n\right|_p < \epsilon$ for $\forall j$. Then $\lim_{j\to\infty} A_j(s) = A_0(s)$ for all $s \in D$. This is used by Washington [24] to show that each of the function $w^{-s}(a)a^s$ and $\sum_{m=0}^{\infty}\binom{s}{m}\left(\frac{F}{a}\right)^m B_m$, where F is the multiple of q and $f = f_\chi$, is analytic in D.

Lemma 3, cf. [24]. *(Washington) Let χ be the primitive Dirichlet character, and let F be a positive integral multiple of q and $f = f_\chi$. Define*

$$L_p(s,\chi) = \frac{1}{s-1}\frac{1}{F}\sum_{\substack{a=1\\(a,p)=1}}^{F} \chi(a) < a >^{1-s} \sum_{m=0}^{\infty}\binom{1-s}{m}\left(\frac{F}{a}\right)^m B_m. \tag{17}$$

Note that $L_p(s,\chi)$ is analytic for $s \in D$ when $\chi \neq 1$, and meromorphic for $s \in D$, with simple pole at $s = 1$ having residue $1 - \frac{1}{p}$ when $\chi = 1$. Furthermore, for each $n \in \mathbb{Z}$, $n \ge 1$, we have

$$L_p(1-n,\chi) = -\frac{1}{n}\left(1 - \chi_n(p)p^{n-1}\right)B_{n,\chi_n}. \tag{18}$$

We now construct the multivariate p-adic L-function, $L_{p,r}(s,\chi)$, which interpolates the multiple generalized Bernoulli numbers associated with χ at negative integers.

Let F be a positive integral multiple of q and $f = f_\chi$, and let us define the multivariate p-adic L-function as follows:
(19)

$$L_{p,r}(s,\chi) = \frac{1}{\prod_{j=1}^{r}(s-j)} \frac{1}{F^r} \sum_{\substack{a_1,\cdots,a_r=1 \\ (a_1+\cdots+a_r,p)=1}}^{F} \chi(a_1+\cdots+a_r) < a_1+\cdots+a_r >^{r-s} \cdot \sum_{m=0}^{\infty} \binom{r-s}{m} \left(\frac{F}{a_1+\cdots+a_r}\right)^m B_m^{(r)}.$$

Then $L_{p,r}(s,\chi)$ is analytic for $t \in \mathbb{C}_p$ with $|t|_p \leq 1$, provided $s \in D$, except $s \neq 1, 2, \cdots, r-1$ when $\chi \neq 1$. We now let $n \in \mathbb{Z}$, $n \geq 1$, and fix $t \in \mathbb{C}_p$ with $|t|_p \leq 1$. Since F must be a multiple of $f = f_{\chi_n}$. By (6), we see that

$$B_{n+r,\chi_{n+r}}^{(r)} = F^n \sum_{a_1,\cdots,a_r=0}^{F-1} B_{n+r}^{(r)} \left(\frac{a_1+\cdots+a_r}{F}\right) \chi_{n+r}(a_1+\cdots+a_r). \tag{20}$$

If $\chi_n(p) = 0$, then $(p, f_{\chi_n}) = 1$, so that $\frac{F}{p}$ is multiple of f_{χ_n}. Let

$$I_0 = \{\frac{a_1+\cdots+a_r}{p} | a_1+\cdots+a_r \equiv 0 (\mod p) \text{ for some } a_i \in \mathbb{Z} \text{ with } 0 \leq a_i \leq F\}.$$

Then we have

$$\begin{aligned} & F^n \sum_{\substack{a_1,\cdots,a_r=0 \\ p|a_1+\cdots a_r}}^{F-1} B_{n+r}^{(r)} \left(\frac{a_1+\cdots+a_r}{F}\right) \chi_{n+r}(a_1+\cdots+a_r) \\ & = p^n \left(\frac{F}{p}\right)^n \chi_n(p) \sum_{\substack{a_1,\cdots,a_r=0 \\ \beta \in I_0}}^{\frac{F}{p}} \chi_{n+r}(\beta) B_{n+r}^{(r)} \left(\frac{\beta}{\frac{F}{p}}\right). \end{aligned} \tag{21}$$

Now we also define the second multiple generalized Bernoulli numbers attached to χ as follows:

$$B_{n+r,\chi_{n+r}}^{*(r)} = \left(\frac{F}{p}\right)^n \sum_{\substack{a_1,\cdots,a_r=0 \\ \beta \in I_0}}^{\frac{F}{p}} \chi_{n+r}(\beta) B_{n+r}^{(r)} \left(\frac{\beta}{\frac{F}{p}}\right). \tag{22}$$

Thus, we note that

$$\begin{aligned} & B^{(r)}_{n+r,\chi_{n+r}} - p^n \chi_n(p) B^{*(r)}_{n+r,\chi_{n+r}} \\ & = F^n \sum_{\substack{a_1,\cdots,a_r=1 \\ p\nmid a_1+\cdots+a_r}}^{F} \chi_{n+r}(a_1+\cdots+a_r) B^{(r)}_{n+r}\left(\frac{a_1+\cdots+a_r}{F}\right). \end{aligned} \tag{23}$$

By the definition of the multiple Bernoulli polynomials of order r, we see that

$$B^{(r)}_{n+r}\left(\frac{a_1+\cdots+a_r}{F}\right) = F^{-n-r}(a_1+\cdots+a_r)^{n+r} \sum_{k=0}^{n} \binom{n+r}{k} \left(\frac{F}{a_1+\cdots+a_r}\right)^k B^{(r)}_k. \tag{24}$$

From (23) and (24), we can derive the below Eq.(25):

$$\begin{aligned} B^{(r)}_{n+r,\chi_{n+r}} - p^n \chi_n(p) B^{*(r)}_{n+r,\chi_{n+r}} = & F^{-r} \sum_{\substack{a_1,\cdots,a_r=1 \\ p\nmid a_1+\cdots+a_r}}^{F} (a_1+\cdots+a_r)^{n+r} \chi_{n+r}(a_1+\cdots+a_r) \\ & \cdot \sum_{k=0}^{n} \binom{n+r}{k} \left(\frac{F}{a_1+\cdots+a_r}\right)^k B^{(r)}_k. \end{aligned} \tag{25}$$

By (18) and (25), we easily see that

$$\begin{aligned} & L_{p,r}(-n,\chi) \\ & = \frac{(-1)^r}{\prod_{j=1}^{r}(n+j)} \frac{1}{F^r} \sum_{\substack{a_1,\cdots,a_r=1 \\ (a_1+\cdots+a_r,p)=1}}^{F} \chi_{n+r}(a_1+\cdots+a_r)(a_1+\cdots+a_r)^{n+r} \\ & \cdot \sum_{m=0}^{\infty} \binom{r+m}{m} \left(\frac{F}{a_1+\cdots+a_r}\right)^m B^{(r)}_m \\ & = (-1)^r \frac{n!}{(n+r)!} \left(B^{(r)}_{n+r,\chi_{n+r}} - \chi_{n+r}(p) p^n B^{*(r)}_{n+r,\chi_{n+r}} \right). \end{aligned}$$

Therefore we obtain the below theorem:

Theorem 4. *Let F be a positive integral of q and $f = f_\chi$, and let*

$$
\begin{aligned}
&L_{p,r}(s,\chi)\\
&= \frac{1}{\prod_{j=1}^{r}(s-j)} \frac{1}{F^r} \sum_{\substack{a_1,\cdots,a_r=1\\(a_1+\cdots+a_r,p)=1}}^{F} \chi(a_1+\cdots+a_r) < a_1+\cdots+a_r >^{r-s}\\
&\cdot \sum_{m=0}^{\infty} \binom{r-s}{m} \left(\frac{F}{a_1+\cdots+a_r}\right)^m B_m^{(r)}.
\end{aligned}
$$

Then, $L_{p,r}(s,\chi)$ is analytic for $t \in \mathbb{C}_p$, $|t|_p \le 1$, provided $s \in D$, except $s \neq 1, 2, \cdots, r$. Also, if $t \in \mathbb{C}_p$, $|t|_p \le 1$, this function is analytic for $s \in D$ when $\chi \neq 1$, and meromorphic for $s \in D$. Furthermore, for each $n \in \mathbb{Z}$, $n \ge 1$, we have

$$
L_{p,r}(-n,\chi) = (-1)^r \frac{n!}{(n+r)!} \left(B_{n+r,\chi_{n+r}}^{(r)} - \chi_{n+r}(p) p^n B_{n+r,\chi_{n+r}}^{*(r)} \right).
$$

The question to construct multivariate p-adic L-function in p-adic number field is still open. Theorem 4 can be considered as a part of answer for the question which was remained in [8].

REFERENCES

1. L. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J. **15** (1948), 987-1000.
2. M. Cenkci, M. Can, V. Kurt, *p-adic interpolation functions and Kummer type congruences for q-twisted and q-generalized twisted Euler numbers*, Advan. Stud. Contemp. Math. **9** (2004), 203-216.
3. G. Choi, A. Zaharescu, *Numerical polynomials in several variables with few nonzero terms*, Advan. Stud. Contemp. Math. **10** (2005), 7-13.
4. H.-S. Cho, E.-S. Kim, *Translation-invariant p-adic integral on $\mathbb{Z}_p$*, Notes on Number Theory and Discrete Mathematics **7** (2001), 70-77.
5. J. Diamond, *The p-adic* log *gamma function and p-adic Euler constant*, Trans. Amer. Math. Soc. **233** (1977), 321-337.
6. B. Ferrero, R. Greenberg, *On the behavior of p-adic L-functions at $s = 0$*, Invent. Math. **50** (1978), 91-102.

7. K. Iwasawa, *Lectures on p-adic L-functions*, Princeton Univ. Press, 1972.

8. T. Kim, *A note on multiple L-function*, Tamsui Oxford J. Math. Sci. (2005), 00-00.

9. T. Kim, *Some formulae for the q-Bernoulli and Euler polynomials of higher order*, J. Math. Anal. Appl. **273** (2002), 236-242.

10. T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288-299.

11. T. Kim, *An invariant p-adic integral associated with Daehee numbers*, Integral Trans. Special Funct. **13** (2002), 65-69.

12. T. Kim, L. C. Jang, S. H. Rim, *An extension of q-zeta function*, Int. J. Math. Math. Sci. **2004 no. 49** (2004), 2649-2651.

13. T. Kim, *On p-adic q-L-functions and sums of powers*, Discrete Math. **252** (2002), 179-187.

14. T. Kim, *On the twisted q-zeta functions and q-Bernoulli polynomials*, Far East J. Appl. Math. **13** (2003), 13-21.

15. T. Kim, *p-adic q-integrals associated with Changhee-Barnes' q-Bernoulli polynomials*, Integral Trans. Special Funct. **15** (2004), 415-420.

16. T. Kim, *Analytic continuation of multiple q-zeta functions and their values at negative integers*, Russian J. Math. Phys. **11** (2004), 71-76.

17. T. Kim, *Non-archimedean q-integrals associated with multiple Changhee q-Bernoulli polynomials*, Russian J. Math. Phys. **10** (2003), 91-98.

18. T. Kim, *A new approach to q-zeta function*, arXiv:math.NT/0502005 (1 Feb 2005).

19. T. Kim, C. Adiga, *Sums of products of generalized Bernoulli numbers*, Inter. Math. J. **5 no. 1** (2004), 1-7.

20. N. Koblitz, *A new proof of certain formulas for p-adic L-function*, Duke Math. J **46** (1979), 455-468.

21. A. A. Panchishkin, *Two variable p-adic L-functions attached to eigenfamiles of positive slope*, Invent. Math. **154** (2003), 551-615.

22. M. Schlosser, *q-analogues of the sums of consecutive integers, squares, cubes, quarts, and quints*, The Electronic J. Combinatorics **11** (2004), R 71.

23. K. Shiratani, S. Yamamoto, *On a p-adic interpolation function for the Euler numbers and its derivatives*, Mem. Fac. Sci.Kyushu Univ. **39** (1985), 113-125.

24. L. C. Washington, *Introduction to cyclotomic fields*, Springer-Verlag(1'st Ed.), 1982.